

- [0594] 13. C. E. Chiche-Lapierre, C. E., Tramalloni, D., Chaput, N. et al. Comparative Analysis of Sepax S-100, COBE 2991, and Manual DMSO Removal Techniques From Cryopreserved Hematopoietic Stem Cell Apheresis Product *Cytotherapy* 2016 18, 6:S47.
- [0595] 14. Huang, L., Cox, E., Austin, R. Continuous particle separation through deterministic lateral displacement, *Science* 2004, 304:987-990.
- [0596] 15. Davis, J. A., Inglis, D. W., et al. Deterministic Hydrodynamics: Taking Blood Apart. *Proc. Natl. Acad. Sci. USA* 2006, 103, 14779-14784.
- [0597] 16. Inglis, D. W., Davis, J. A., Austin, R. H. Critical particle Size for Fractionation by Deterministic Lateral Displacement. *Lab Chip* 2006, 6, 655-658. 17. Chen, Y., D'Silva, J. Austin, R. et al. Microfluidic chemical processing with on-chip washing by deterministic lateral displacement arrays with separator walls. *Biomicrofluidics*. 2015 9(5): 054105.
- [0598] 18. Shilun, F. Skelley, A., Anwer, A. G., et al. Maximizing Particle Concentration in Deterministic Lateral Displacement Arrays. *Biomicrofluidics* 2017, 11, 024121.
- [0599] 19. D'Silva, J. Throughout Microfluidic Capture of Rare Cells from Large Volumes of Blood. A Dissertation Presented to the Faculty of Princeton University in Candidacy for the Degree of Doctor of Philosophy. 2016.
- [0600] 20. NIST/SEMATECH e-Handbook of Statistical Methods, 2017, www.itl.nist.gov/div898/handbook/August
- [0601] 21. Mahnke, Y. D., Brodie, T. M., Sallusto, F., et al. The Who's Who of T-cell Differentiation: Human Memory T-cell Subsets. *Eur. J. of Immunol.* 2013, 43, 2797-2809.
- [0602] 22. Civin, C. I., Ward, T., Skelley, A. M., et al. Automated Leukocyte Processing by Microfluidic Deterministic Lateral Displacement. *Cytometry A*. 2016, 89, 1073-1083.
- [0603] 23. Trickett, A., and Kwan, Y. L. T-cell Stimulation and Expansion Using Anti-CD3/CD28 Beads. *J. Immunol. Meth.* 2003, 275, 251-255.
- [0604] 24. Marktkamcham, S., Onlamoon, N., Wang, S., et al. The Effects of Anti-CD3/CD28 Coated Beads and IL-2 on Expanded T-cell for Immunotherapy. *Adv. Clin. Exp. Med.* 2016, 25, 821-828.
- [0605] 25. Li, Y., and Kurlander, R. J. Comparison of Anti-CD3 and Anti-CD28-coated Beads with Soluble Anti-CD3 for Expanding Human T-cells: Differing Impact on CD8 T-cell Phenotype and Responsiveness. *J. Transl. Med.* 2010, 8, 104-118.
- [0606] 26. Agrawal S., Ganguly, S., Hjian, P., et al. PDGF Upregulates CLEC-2 to Induce T Regulatory Cells. *Oncotarget*. 2015, 6, 28621-28632.
- [0607] 27. Zhu, L., Huang, Z., Stålesen, R. et al. Platelets Provoke Distinct Dynamics of Immune Response by Differentially Regulating CD4⁺ T-cell Proliferation. *J. Throm. Haem.* 2014, 12, 1156-1165.
- [0608] 28. Koesdjojo, M., Lee, Z., Dosier, C., et al. DLD Microfluidic Purification and Characterization of Intact and Viable Circulating Tumor Cells in Peripheral Blood. *AACR Annual Meeting* 2016, Abstract #3956.
- [0609] 29. Loutherbak, K. "Microfluidic Devices for High Throughput Cell Sorting and Chemical Treatment," A Dissertation Presented to the Faculty of Princeton University 2011.
- [0610] All references cited herein are fully incorporated by reference. Having now fully described the invention, it will be understood by one of skill in the art that the invention may be performed within a wide and equivalent range of conditions, parameters and the like, without affecting the spirit or scope of the invention or any embodiment thereof
- What is claimed is:
1. A method for preparing cells for treating a patient with cancer, comprising the steps:
 - a) purifying T cells from a sample comprising leukocytes and platelets, wherein the leukocytes comprise T cells and wherein the T cells are purified by:
 - i) performing a size based separation using a microfluidic device to produce an enriched product in which, compared to the sample, the percentage cells that are platelets has been reduced; and
 - ii) in addition to the size based separation, performing an affinity based separation;
 - b) after the purification of step a), activating and expanding the T cells to produce a composition in which the percentage of T cells that are central memory T cells has increased compared to the percentage of T cells that are central memory T cells in the sample;
 - c) genetically engineering activated T cells to comprise therapeutic benefit in the treatment of said patient's cancer.
 2. The method of claim 1, wherein, in step c), the activated T cells are genetically engineered to comprise modified cell surface receptors of therapeutic benefit in the treatment of said patient's cancer.
 3. The method of claim 2, wherein, the modified cell surface receptors of therapeutic benefit are chimeric antigen receptors (CARs).
 4. The method of claim 1, wherein the sample is obtained by apheresis
 5. The method of claim 1, wherein the sample is obtained by leukapheresis.
 6. The method of claim 1, wherein the platelets in the enriched product of paragraph a)ii) are depleted by at least 80% compared to the sample and/or there are no more than 5 platelets per leukocyte in the enriched product.
 7. The method of claim 1, wherein, the genetically engineered T cells are collected by transferring them into a pharmaceutical composition for administration to a patient.
 8. The method of claim 7, wherein cells are not frozen before being collected.
 9. A method of treating a patient for cancer comprising administering to said patient genetically engineered T cells prepared by a method comprising the steps of:
 - a) purifying T cells from a sample comprising leukocytes and platelets, wherein the leukocytes comprise T cells and wherein the T cells are purified by:
 - i) performing a size based separation using a microfluidic device to produce an enriched product in which, compared to the sample, the percentage cells that are platelets has been reduced; and
 - ii) in addition to the size based separation, performing an affinity based separation;
 - b) after the purification of step a), activating and expanding the T cells to produce a composition in which the percentage of T cells that are central memory T cells has increased compared to the percentage of T cells that are central memory T cells in the sample;